

UDC: 576.316.352: 595.773.4:539.1.047

Особливості гомологічної кон'югації політенних хромосом за впливу мікрохвиль і статичного магнітного поля у *Drosophila melanogaster* Meig.
Л.Д.Дика, Л.О.Шакина, В.Ю.Страшнюк

Харківський національний університет імені В.Н.Каразіна (Харків, Україна)
lilya_dikaya@ukr.net, LubovZ2003@gmail.com, volodymyr.strashnyuk@gmail.com

Метою роботи було вивчити вплив мікрохвильового випромінювання, статичного магнітного поля та їх сумісної дії на порушення гомологічного спарювання політенних хромосом у *Drosophila melanogaster* Meig. Матеріалом для дослідження була інbredна лінія *Canton-S*. Застосовували мікрохвилі з частотою 36,64 ГГц і щільністю потужності 1 Вт/м² протягом 30 секунд і статичне магнітне поле напруженістю 25 мТл протягом 5 хв. Вплив електромагнітних полів здійснювали на стадії яйця після 2-годинної яйцекладки. Політенні хромосоми досліджували на давлених препаратах слинних залоз, забарвлених ацетоорсеїном. Для приготування препаратів використовували самок на стадії 0-годинної передлялечки. Мікрохвилі зменшували частоту асинапсису гомологічних хромосом на 18,1%. Статичне магнітне поле не мало значущого впливу. Комбінована дія цих чинників зменшувала частоту асинапсису в гігантських хромосомах дрозофіли на 37,1%. Отримані результати свідчать про вплив електромагнітних чинників на гомологічну кон'югацію хромосом в ядрах соматичних клітин. Таким чином, показана можливість їх впливу на транс-взаємодії локусів гомологічних хромосом.

Ключові слова: гігантські хромосоми, дрозофіла, гомологічна кон'югація, мікрохвилі, статичне магнітне поле, транс-взаємодії.

Peculiarities of homologous conjugation of polytene chromosomes after exposure to microwaves and static magnetic field in *Drosophila melanogaster* Meig.

L.D.Dyka, L.A.Shakina, V.Yu.Strashnyuk

The purpose of this investigation was to study the effect of microwave irradiation, static magnetic field and their combined action on the violation of homologous pairing of polytene chromosomes in *Drosophila melanogaster* Meig. Inbred line *Canton-S* was used as an experimental object. Microwaves with frequency 36.64 GHz and power density 1 W/m² for 30 sec and static magnetic field of 25 mT intensity for 5 min were used. Exposure to electromagnetic fields was applied at the egg stage after 2-hour oviposition. Polytene chromosomes were examined in squashed salivary glands preparations stained with acetoorcein. For preparations, females at the 0-hour prepupa stage were used. Microwaves decreased the asynapsis frequency of homologous chromosomes by 18.1%. The constant magnetic field had no significant effect. Combined action of these factors decreased the asynapsis frequency in *Drosophila* giant chromosomes by 37.1%. Obtained results testify to the influence of electromagnetic factors on the homologous pairing of chromosomes in the nuclei of somatic cells. Thus, the possibility of their effect on the trans-interactions of the loci of homologous chromosomes is shown.

Key words: giant chromosomes, *Drosophila*, homologous conjugation, microwaves, static magnetic field, trans-interactions.

Особенности гомологичной конъюгации политенных хромосом при действии микроволн и постоянного магнитного поля у *Drosophila melanogaster* Meig.

Л.Д.Дикая, Л.А.Шакина, В.Ю.Страшнюк

Целью работы было изучить влияние микроволнового излучения, постоянного магнитного поля и их сочетанного действия на нарушение гомологичного спаривания политенных хромосом у *Drosophila melanogaster* Meig. Материалом для исследования служила инbredная линия *Canton-S*. Применяли микроволны с частотой 36,64 ГГц и плотностью мощности 1 Вт/м² в течение 30 секунд и постоянное магнитное поле напряжённостью 25 мТл в течение 5 мин. Воздействие электромагнитных полей производили на стадии яйца после 2-часовой яйцекладки. Политенные хромосоми исследовали на давлених препаратах слюнных желёз, окрашенных ацетоорсеїном. Для приготовления препаратів використовували самок на стадії 0-часової предкуколки. Микроволны уменьшали частоту асинапсису гомологічних хромосом на 18,1%. Постоянное магнитное поле не имело значимого эффекта.

Комбинированное действие этих факторов уменьшало частоту асинопсиса в гигантских хромосомах дрозофилы на 37,1%. Полученные результаты свидетельствуют о влиянии электромагнитных факторов на гомологичную конъюгацию хромосом в ядрах соматических клеток. Таким образом, показана возможность их влияния на транс-взаимодействия локусов гомологичных хромосом.

Ключевые слова: гигантские хромосомы, дрозофила, гомологичная конъюгация, микроволны, постоянное магнитное поле, транс-взаимодействия.

Introduction

In recent decades a new important environmental factor has emerged and developed – electromagnetic fields (EMFs) of anthropogenic origin (Shckorbatov, 2014). Technological equipments for various purposes using microwave radiation are widely introduced. These are variable and pulsed magnetic fields, medical therapeutic and diagnostic facilities, visual information displays on cathode ray tubes (monitors, computers, televisions, etc.), industrial equipment with power supply, household appliances, personal communications (mobile phones) and the like. For today, we can state an increase in the effect of electromagnetic radiation on living organisms. The World Health Organization has included this issue in the list of the most pressing for humanity (WHO, 1997). International Agency for Research on Cancer (IARC) considers radiofrequency EMFs as potentially carcinogenic to humans (WHO/IARC, 2011).

The biological effects of EMFs are intensively studied at various objects and at different levels of the organization. In view of genetics, research on the chromosome level is of particular interest. A convenient object for this is the giant chromosomes of Diptera, which represent a modification of the actively functioning interphase chromosomes.

The capacity for homologous pairing is a fundamental property of eukaryotic chromosomes. Homologous and non-homologous (ectopic) chromosome associations play an important role in the spatial organization of the cell nucleus, providing functional connections of the different elements of genome. This issue is being considered currently in the light of the epigenetic control of gene expression (Duncan, 2002; Stegny, 2006; Schneider, Grosschedl, 2007).

In most cases, homologous chromosomes in *Drosophila* polytene nuclei tightly conjugated to each other. But often the conjugation is broken, there is an asynapsis phenomenon.

Asynapsis was investigated in homo- and heterozygous organisms. The interline differences in the frequency of violations of homologous chromosomes pairing were shown. The increase of spontaneous asynapsis was observed due to hybridization in the study of heterosis phenomenon (Lapta, Shakhbazov, 1976; Shakina et al., 2005; Taglina, 2006), as well as in interspecific hybrids of *Drosophila* (Evgen'ev, Polianskaya, 1976).

There are too few works, dealing with violations of conjugation of polytene chromosomes due to exogenous influences (Navrotskaya, 2006; Vasserlauf et al., 2008). Researches in this direction are very promising, given the association of terminal points of asynapsis with heterochromatic regions (Lapta, 1977), which are known to have a high polymorphism and dynamic structure (Grewal, Jia, 2007).

It should be also noted a lack of researches on the effects of EMFs on the chromosomal level (Tonomura et al., 1990, 1992; Goodman, Blank, 1998). In our previous works, we studied the effect of microwaves irradiation and a constant magnetic field on the puffing activity (Shakina et al., 2011) and endoreduplication (Dyka et al., 2016) in polytene chromosomes. An additional reason for our study were modifications of trans-inactivation effect under the influence of microwave irradiation, depending on the presence or absence of synapsis of homologous chromosomes, found in experiments on *Drosophila* (Shckorbatov et al., 2004).

The purpose of this investigation was to study the effects of microwave radiation, a constant magnetic field and their combined action on violations of the homologous pairing of polytene chromosomes in *Drosophila melanogaster* Meig.

Objects and methods of research

Studies were carried out on inbred strain *Canton-S*. This strain was obtained from *Drosophila* collection of the Department of Genetics and Cytology of the Kharkiv National University through inbred breeding by mating siblings. The degree of inbreeding before experiments was 77 generations.

Flies were grown on standard sugar-yeast medium at $24.0 \pm 0.5^\circ\text{C}$. *Drosophila* culture developed in vials of 60 ml. The volume of culture medium in each vial was 10 ml.

EMF exposures were done by using an microwave emitter, which generates linearly polarized EMF with a frequency of 36.64 ± 0.05 GHz. This semi-conductor device was designed and constructed at the

Department of Theoretical Radiophysics of V.N.Karazin Kharkiv National University based Gunn diode (the author is V.N.Bykov). Microwaves of this frequency refer to the Ka band (27–40 GHz). Such radiation is used in various radar systems (Richards, 2014). The power density in our experiments was 1 W/m^2 , exposure time – 30 sec. The distance between the emitting antenna horn edge and the eggs was 15 cm. The SAR (Specific Absorption Rate) was 4 W/kg . In the literature, the level of microwave radiation below 100 W/m^2 assumed as not thermal (Michaelson, 1980).

In addition, exposure to static magnetic field with 25 mT intensity for 5 min was applied. We studied separately the effects of a single exposure to microwaves or magnetic field as well as the sequential action of microwaves and magnetic fields.

Exposure to EMFs was applied in early embryogenesis. We used newly-laid *Drosophila* eggs obtained from the five-day-old flies after a short 2-hour lay. Eggs were laid on the fresh culture medium surface and then were exposed. Examinations of the effects of EMFs were performed at the 0-hour prepupa stage. It is known that asynapsis pattern is formed in the early stages of development and doesn't vary in larvae of different age (Lapta, Shakhbazov, 1976). In the control experiments *Drosophila* "sham eggs" were not exposed to microwave irradiation and magnetic field.




Fig. 1. Giant chromosomes of *Drosophila melanogaster* stained by aceto-orcein with different number of asynapsis regions: (a) asynapsis is absent; (b) a single asynapsis; (c) two asynapsis segments; (d) three asynapsis segments

Polytene chromosomes were examined on squashed salivary glands preparations, stained with aceto-orcein: 2% orcein (Merck KGaA, Darmstadt, Germany) solution in 45% acetic acid (Reahimtrans, Kyiv, Ukraine). For preparations, females at the 0 hours prepupae stage were used. The preparations were analyzed by light microscopy (MBI-6 microscope, "LOMO", St. Petersburg, Russia) at magnification $\times 200$. We assessed the percentage of nuclei with different number of asynapsis regions in the preparations: no asynapsis; a single asynapsis segment; two asynapsis segments; three and more asynapsis segments (Fig. 1). The asynapsis frequency was defined as the ratio of non-conjugating chromosomal segments to the total number of investigated nuclei.

Experimental data were statistically analysed. Data are presented as the mean \pm standart error. For each experiment 10 larvae were used. The asynapsis regions were determined in 25–50 nuclei per preparation. We examined from 297 to 392 nuclei in each variant of experiment.

The significance of differences between the study groups was assessed using Student's *t*-test with the Bonferroni amendment for multiple comparisons. Results considered valid when $p < 0.017$.

Results


The percentage of nuclei with different number of asynapsis segments in *Drosophila melanogaster* in the control and after exposure to electromagnetic factors are presented in Fig. 2.

In the control group violations of conjugation of homologous chromosomes weren't found in 67.3% of the examined nuclei. The single asynapsis was revealed in 21.7% of the nuclei, 9.2% of nuclei had two violations of conjugation, in 1.8% cases three or more asynapsis segments were found.

After the exposure to microwave radiation, as well as after the action of a constant magnetic field, the percentage of nuclei without asynapsis didn't differ from control values. After the microwave irradiation the portion of nuclei with a single asynapsis section increased by 22.6%, the number of nuclei with two violations of synapsis decreased by 55.4%, while the percentage of nuclei with three or more asynapsis segments didn't differ from the control.

After the action of a constant magnetic field the content of nuclei with a single, as well as three or more violations of synapsis was the same as in the control, the proportion of nuclei with two asynapsis decreased by 39.1%.


The most significant changes were found after the combined effects of microwaves and static magnetic field. The portion of nuclei without violations of chromosomes' pairing increased by 11.4%. The number of nuclei with a single asynapsis segment wasn't changed, the portion of nuclei with two asynapsis regions decreased by 62.0%, while the nuclei with three or more violations of synapsis weren't found.



* – $p < 0,017$; ** – $p < 0,01$; *** – $p < 0,00$: versus to control group

Fig. 2. The ratio of nuclei with different numbers of asynapsis segments in polytene chromosomes after microwave irradiation, the action of a constant magnetic field and their combined influence in *Drosophila melanogaster*: 1 – asynapsis is absent; 2 – a single asynapsis; 3 – two asynapsis segments; 4 – three asynapsis segments

The data on the portion of nuclei with different numbers of asynapsis segments were used for calculation of average frequency of violations of homologous conjugation in giant chromosomes in *Drosophila melanogaster* in norm and after experimental influences. The results are presented in Fig. 3.



* – $p < 0,017$; *** – $p < 0,001$: versus to control group

Fig. 3. The frequency of violations of homologous conjugation in polytene chromosomes of *Drosophila melanogaster* after the microwave irradiation, the action of a constant magnetic field and their combined effect

The average frequency of asynapsis in the control group was 45.2%. After the microwave irradiation decrease of the asynapsis frequency by 18.1% was observed. The influence of constant magnetic field wasn't statistically significant. The largest effect occurred when combined action of microwave radiation and magnetic field was applied: frequency of asynapsis decreased by 37.4% compared with the control. This index was also lower than after exposure to microwaves – by 23.5%, or a constant magnetic field – by 30.6%.

Thus, the 36.64 GHz microwave irradiation on the stage of embryogenesis with power density 1 W/m² and 30 sec of exposure led to reduction in of the homologous chromosomes asynapsis frequency in the salivary glands cells in *Drosophila melanogaster*. The constant magnetic field with 25 mT intensity for 5 min hadn't significant effect on the asynapsis frequency of itself, but significantly increased the effect of microwave radiation.

Discussion

The violations of conjugation of homologous chromosomes are of two types: specific and non-specific, or spontaneous. The first type is a regular asynapsis and associated with various pattern of chromosomal discs. This occurs, for example, as a result of interspecific hybridization (Evgen'ev, Polianskaya, 1976). The second type occurs when the picture of disks is identical and its nature is not fully understood (Lapta, 1977; Taglina, 2006).

There is ample evidence, which indicates that the principles of somatic and meiotic conjugation are the same (Martinez-Perez et al., 2001). In the literature, the synapsis of homologous chromosomes is explained by preferential attraction of certain chromosomal regions, which localization usually correlates with position of heterochromatin (Tiang, Pawlowski, 2012). It has been also shown that zones of violation of conjugation begin or end in the centromeric, telomeric and intercalary heterochromatin (Lapta, 1977).

The character of homologues interaction affects in a certain way on gene expression. There are examples of puffs heterozygosity in sites of asynapsis in interlineal hybrids of *Drosophila*: sizes of puffs derived from different parental lines differ in the asinapsys segments of homologous chromosomes (Ashburner, 1970; Strashnyuk et al., 2009). Such examples are also characteristic for interspecific hybrids in which the regularly asynapsis occurs apart from the spontaneous one (Ashburner, 1969).

Ashburner (1970) showed the influence of presence or absence of physical contact of homologues chromosomes on gene expression. In his view, synapsis may correct damages leading to restoration of gene activity that affects the puff morphology.

At present, it has been proven possible to restore a normal phenotype by interaction of differently

damaged genes that are located in homologous chromosomes when they physically contact. This phenomenon is called transvection. Broadly, the transvection is such cis-trans interactions, which may be accompanied by either activation or inactivation of genes (Henikoff et al., 1995; Kennison, Southworth, 2002; Csink et al., 2002). This phenomenon is associated with subtle mechanisms of genes activity regulation observed in eukaryotic organisms of different taxa: fungi, higher plants, insects, mammals (Duncan, 2002).

An example is the effect of trans-inactivation (Kennison, Southworth, 2002; Csink et al., 2002). To study this effect, in particular, *Drosophila* line, comprising a mutation *brown*^{Dominant} (*bw^D*), is used. *bw^D* gene is able to inhibit the expression of the gene *bw^t* and shows the effect of dominance in the case, if the conjugation of chromosomes occurs in the area in which *bw^D* and *bw^t* genes are localized (Henikoff et al., 1995). Thus, by examining the expression of these genes, it is possible to judge on the changes in the subtle processes of chromosome interaction inside the cell nucleus.

Earlier in the line containing the mutation *bw^D*, the effect of microwave radiation on the manifestation of trans-inactivation effect in *Drosophila* was studied (Shckorbatov et al., 2004). The linearly polarized and the left polarized radiation with $\lambda=8$ mm, $E=0.2$ mW/cm² caused increased trans-inactivation effect, manifested in the reduction of pigment in the eyes of the flies. This indicated an increase in homologous chromosomes conjugation degree in the cell nucleus. Right polarized radiation, on the contrary, caused an increase in pigment content, i.e. trans-inactivation effect decreased. Thus, it was shown the effect of electromagnetic waves on gene expression, which is associated with the state of conjugation of chromosomes in the nucleus.

The data of this study are consistent with the effect of microwaves on the manifestation of trans-inactivation effect. In our study, we applied linearly polarized EMF. The frequency of violations of homologous chromosomes pairing reduced as in the study of trans-inactivation effect. In the case of trans-inactivation authors judged about homologous chromosomes interaction indirectly, based on the data on mutant trait expressivity changes. The study of polytene chromosomes allows us to evaluate it directly by visual observation.

The effect of electromagnetic fields on the interaction of homologous chromosomes can be explained by taking into account the data about their impact on chromatin structure. Earlier in the human cells, it has been shown an increase of heterochromatin granules content in the nuclei under the influence of microwaves (Shckorbatov et al., 2009). Given that heterochromatic regions are the sites of synapsis initiation, the data on the increase of genome heterochromatinization degree under the influence of electromagnetic fields is consistent with homologous chromosomes asynapsis decrease. The dependence of asynapsis frequency on the amount of heterochromatin was demonstrated earlier (Belyaeva, 1973; Lapta, 1977).

Heterochromatinization apparently is a manifestation of a more general cell response on the stress, as DNA in a tightly packed chromatin is better protected against damaging influences. The signal for the conformational changes in the chromatin may be due to the change of ionic composition in cytoplasm and karyoplasm associated with the changes in membrane permeability, which takes place under microwave irradiation of the cell (Skamrova et al., 2013).

The magnetic fields also can induce the stress response in Diptera (Goodman, Blank, 1998). Exposure to low frequency magnetic field induced transcription in specific regions of giant chromosomes in *Drosophila melanogaster* and *Sciara coprophila*. The activation of heat shock transcriptional factor (HSF1), increase in the synthesis of stress protein hsp 70, and 3–4-fold increase of heat shock element binding activity were shown.

Conclusions

In the present study, exposure to microwave radiation (frequency – 36.64 GHz, power density – 1 W/m², exposure time – 30 sec) on the stage of embryogenesis decreased the asynapsis frequency of homologous chromosomes in the salivary glands cells of *Drosophila melanogaster* by 18.1%. The constant magnetic field (intensity 25 mT, exposure time – 5 min) had no significant effect. Combined action of microwaves and static magnetic field decreased the asynapsis frequency in *Drosophila* polytene chromosomes by 37.1%. Obtained results testify to the influence of electromagnetic factors on the homologous pairing of chromosomes in the nuclei of somatic cells. Thus, the possibility of their effect on the trans-interactions of the loci of homologous chromosomes is shown.

Acknowledgements

The authors are grateful to Professor L.A.Atramentova, the Head of Department of Genetics and Cytology, V.N.Karazin Kharkiv National University, for helpful advice on statistical methods.

References

- Ashburner M. Patterns of puffing activity in the salivary gland chromosomes of *Drosophila*. III. A comparison of the autosomal puffing patterns of the sibling species of *Drosophila melanogaster* and *Drosophila simulans* // *Chromosoma*. – 1969. – Vol.27. – P. 64–85.
- Ashburner M. The genetic analysis of puffing in polytene chromosomes of *Drosophila* // *Proc. Roy. Soc. Lond.* – 1970. – B176. – P. 319–327.
- Belyaeva E.S. Asynapsis of homologues in *Drosophila melanogaster* salivary chromosomes // *Dros. Inf. Serv.* – 1973. – Vol.50. – P.40.
- Csink A.K., Bounoutas A., Michelle L. et al. Differential gene silencing by *trans*-heterochromatin in *Drosophila melanogaster* // *Genetics*. – 2002. – Vol.160. – P. 257–269.
- Duncan I.W. Transvection effects in *Drosophila* // *Annu. Rev. Genet.* – 2002. – Vol. 36. – P. 521–556.
- Dyka L.D., Shakina L.A., Strashnyuk V.Yu., Shckorbatov Yu.G. Effects of 36,6 GHz and static magnetic field on degree of endoreduplication in *Drosophila melanogaster* polytene chromosomes // *Int. J. Radiat. Biol.* – 2016. – Vol.92 (4). – P. 222–227.
- Evgen'ev M.B., Polianskaya G.G. The pattern of polytene chromosome synapsis in *Drosophila* species and interspecific hybrids // *Chromosoma*. – 1976. – Vol.57. – P. 285–295.
- Goodman R., Blank M. Magnetic field stress induces expression of hsp 70 // *Cell Stress Chaperon*. – 1998. – Vol.3 (20). – P. 79–88.
- Grewal S.I.S., Jia S. Heterochromatin revisited // *Nat. Rev. Genet.* – 2007. – Vol.8. – P. 35–46.
- Henikoff S., Jackson J.M., Talbert P.B. Distance and pairing effects on the brown Dominant heterochromatic element in *Drosophila* // *Genetics*. – 1995. – Vol.140. – P. 1007–1017.
- Kennison J.A., Southworth J.W. Transvection in *Drosophila* // *Adv. Genet.* – 2002. – Vol.46. – P. 399–420.
- Lapta G.E. Localization of asynaptic segments for 2R and 3L chromosomes with a failure of pairing of polytene chromosomes in *Drosophila melanogaster* // *Genetics (USSR)*. – 1977. – Vol.13 (6). – P. 1064–1072. (in Russian)
- Lapta G.E., Shakhbazov V.G. The pattern of asynapsis of polytene chromosomes in inbred lines of *Drosophila melanogaster* and in the F₁ derived from reciprocal crosses between them // *Genetics (USSR)*. – 1976. – Vol.12 (2). – P. 121–126. (in Russian)
- Martinez-Perez E., Shaw P., Moore G. The Ph1 locus is needed to ensure specific somatic and meiotic centromere association // *Nature*. – 2001. – Vol.411. – P. 204–207.
- Michaelson S.M. Microwave biological effects: an overview // *Proc. IEEE*. – 1980. – Vol.68 (1). – P. 40–49.
- Navrotskaya V.V. Manifestation of quantitative traits in intrastain crosses depending on parent generation keeping conditions in *Drosophila melanogaster* Meig. and *Bombyx mori* L. Thesis for Candidate of Biological Sciences degree / 03.00.15 – genetics. – V.N.Karazin Kharkiv National University, Kharkiv, 2006. – 20p.
- Richards M.A. Fundamentals of radar signal processing. 2nd ed. – McGraw-Hill Education, 2014. – 656p.
- Shakina L.A., Pasiuga V.N., Dumin O.M., Shckorbatov Yu.G. Effects of microwaves on the puffing pattern of *D. melanogaster* // *Cent. Eur. J. Biol.* – 2011. – Vol.6 (4). – P. 524–530.
- Shakina L.A., Strashnyuk V.Yu., Shakhbazov V.G. Peculiarities of homologous and non-homologous pairing of polytene chromosomes in the inbred lines and hybrids of *Drosophila* // *The Journal of V.N.Karazin National University. Series "Biology"*. – 2005. – Issue 709 (1–2). – P. 105–110. (in Russian)
- Shckorbatov Y. The main approaches of studying the mechanisms of action of artificial electromagnetic fields on cell // *J. Electr. Electron. Syst.* – 2014. – Vol.3 (2). – P.123.
- Shckorbatov Y.G., Evseeva M.V., Shakhbazov V.G. et al. The influence of the microwave radiation of different polarization on transinactivation effect and viability of *Drosophila* // *Bulletin of Problems Biology and Medicine*. – 2004. – Vol.4. – P. 36–41. (in Russian)
- Shckorbatov Yu.G., Pasiuga V.N., Kolchigin N.N. et al. The influence of differently polarised microwave radiation on chromatin in human cells // *Int. J. Radiat. Biol.* – 2009. – Vol.85 (4). – P. 322–329.
- Skamrova G.B., Lantushenko A.O., Shckorbatov Yu.G., Evstigneev M.P. Influence of mobile phone radiation on membrane permeability and chromatin state of human buccal epithelium cells // *Biochemistry and Biophysics*. – 2013. – Vol.1 (2). – P. 22–28.

- Schneider R., Grosschedl R. Dynamics and interplay of nuclear architecture, genome organization, and gene expression // *Gene. Dev.* – 2007. – Vol. 21. – P. 3027–3043.
- Stegniy V.N. Evolutionary significance of chromosome architecture for epigenetic control of eukaryote development and phylogeny // *Russ. J. Genet.* – 2006. – Vol.42 (9). – P. 1215–1224. (in Russian)
- Strashnyuk V.Yu., Taglina O.V., Gorenskaya O.V., Shakina L.A. Some peculiarities of structural and functional organization of polytene chromosomes of *Drosophila melanogaster* in relation to heterosis // *Factors in experimental evolution of organisms.* – Kyiv: Logos, 2009. – Vol.6. – P. 187–191. (in Russian)
- Taglina O.V. Investigation of spontaneous asynapsis of salivary gland polytene chromosomes of *Drosophila melanogaster* in highly inbred lines and their hybrids // *The Journal of V.N.Karazin National University. Series "Biology".* – 2006. – Issue 729 (3). – P. 136–140. (in Russian)
- Tiang Ch.-L., Pawlowski W.P. Role of telomeres and centromeres in meiotic chromosome pairing // In: *Access Science*, ©McGraw-Hill Education, 2012. (<http://www.accessscience.com>)
- Tonomura Y., Shima T., Suzuki K., Kishi M. Effects of microwaves and magnetic fields on *Drosophila* // *Sci. Rep. Tokyo Woman's Christ. Univ.* – 1990. – Vol.40. – P. 1031–1049.
- Tonomura Y., Shima T., Kimura K. Effects of microwaves on *Drosophila*. Part 2 // *Rep Tokyo Woman's Christ Univ.* – 1992. – Vol.42. – P. 1159–1175.
- Vasserlauf I.E., Shelkovnikova T.A., Mitrenina E.Y., Stegny V.N. The effect inbreeding and low temperature on the pattern of chromosome synapsis in the ovarian nurse nuclei of *Drosophila melanogaster* strains // *Russ. J. Genet.* – 2008. – Vol.44 (8). – P. 1066–1074. (in Russian)
- WHO/International Agency for Research on Cancer (IARC). 2011. Press Release No 208, 31 May.
- WHO International EMF Project. 1997.

Представлено: Н.О.Мазник / Presented by: N.A.Maznyk
Рецензент: Ю.Г.Шкорбатов / Reviewer: Y.G.Shckorbatov
Подано до редакції / Received: 16.10.2017